

Distinct expression patterns of β -1,3-glucanases and chitinases during the germination of Solanaceous seeds

Luciana Petruzzelli¹, Kerstin Müller², Katrin Hermann² and Gerhard Leubner-Metzger^{2*}

¹Istituto Biosintesi Vegetali, C.N.R., via Bassini 15, I-20133 Milano, Italy; ²Institut für Biologie II, Albert-Ludwigs-Universität, Schänzlestr. 1, D-79104 Freiburg i. Br., Germany

Abstract

The expression patterns of β -1,3-glucanases (β Glu) and chitinases (Chn) were investigated during the seed germination of members of the *Cestroideae* (three *Nicotiana* species, *Petunia hybrida*) and the *Solanoideae* (*Capsicum annuum*, *Physalis peruviana*) subgroups of Solanaceous species. Rupture of the micropylar testa (seed coat) and rupture of the micropylar endosperm, i.e. radicle emergence, were distinct and temporally separate events during the germination of *Cestroideae*-type seeds. β Glu accumulation in imbibed *Cestroideae*-type seeds, occurring after testa rupture but prior to endosperm rupture, was inhibited by abscisic acid (ABA) and promoted by gibberellins (GA) and light, in strict association with germination, and appeared to be caused by transcriptional regulation of the class I β Glu genes. The micropylar cap of *Solanoideae*-type seeds does not allow a distinction between testa and endosperm rupture, but β Glu accumulation occurred prior to radicle emergence of pepper and *P. peruviana* seeds. ABA inhibited endosperm rupture and β Glu accumulation in the micropylar cap of pepper seeds. In contrast to tomato, β Glu accumulation in pepper seeds was not only confined to the micropylar cap, was due to distinct, tissue-specific β Glu isoforms, and was not accompanied by Chn accumulation. In conclusion, ABA inhibition of germination and β Glu accumulation in the micropylar endosperm appears to be a widespread event during the seed germination of Solanaceous species. In contrast, accumulation of Chn and distinct β Glu isoforms in the embryo, prior to germination, appears to be a species-specific phenomenon within the *Solanaceae*. In addition, a post-germination co-induction of β Glu and Chn in the root of the emerged seedling was found in endospermic

and non-endospermic species and could represent an evolutionarily conserved event during dicot seedling development.

Keywords: abscisic acid, *Capsicum*, chitinase, gibberellin, β -1,3-glucanase, *Nicotiana*, seed germination, *Solanaceae* family

Introduction

The intrafamilial relationships of Solanaceous species have been investigated using morphological and molecular criteria. On the morphological level, the *Solanaceae* family can be divided into two large subgroups (Judd *et al.*, 1999): (1) The *Cestroideae* subgroup, e.g. *Nicotiana* and *Petunia*, is characterized by straight to slightly bent embryos and prismatic to subglobose seeds, and typically by capsules; and (2) the *Solanoideae* subgroup, e.g. *Capsicum*, *Lycopersicon* and *Physalis*, is characterized by curved embryos and flattened, discoid seeds and often by berries. Seed germination of tobacco (*Nicotiana tabacum*) and tomato (*Lycopersicon esculentum*), type members for each of the two subgroups, is regulated by the balance of forces between the growth potential of the embryo and the micropylar layers that cover the radicle tip and function as a constraint to radicle protrusion (reviewed by Hilhorst, 1995; Bewley, 1997b; Koornneef *et al.*, 2002; Leubner-Metzger, 2003). These covering layers are the testa (seed coat), an entirely maternal tissue, and the triploid endosperm, a predominantly maternal tissue. Both micropylar layers are involved in controlling tobacco and tomato coat-imposed dormancy (e.g. Hilhorst and Downie, 1996; Nonogaki *et al.*, 2000; Wu *et al.*, 2000; Leubner-Metzger, 2002).

Surgical removal of the micropylar testa and the endosperm tissues permits radicle growth under conditions that inhibit germination of intact seeds of tobacco (Bihlmeier, 1927; Kincaid, 1935) and tomato (Liptay and Schopfer, 1983; Hilhorst, 1995). Microscopic studies in tobacco showed that storage

*Correspondence

Fax: + 49 761 2032612

Email: leubner@uni-freiburg.de

Web site: 'The Seed Biology Place' <http://www.leubner.ch>

Abbreviations: ABA = abscisic acid; Chn = chitinase; Chn I = class I chitinase; GA = gibberellin; β Glu = β -1,3-glucanase; β Glu I = class I β -1,3-glucanase.

reserves are degraded in the micropylar endosperm cells prior to protrusion by the radicle and that the endospermic hole, which has a smooth outline and is always formed at the micropylar end of germinating tobacco seeds, results from tissue dissolution rather than from the pushing action of the protruding radicle (Arcila and Mohapatra, 1983; Leubner-Metzger, 2002). Similar observations are obvious for other Solanaceous species and are correlated with weakening of the micropylar endosperm, e.g. in tomato (Hilhorst, 1995; Bewley, 1997a; Toorop *et al.*, 2000) and in pepper (Watkins and Cantliffe, 1983; Watkins *et al.*, 1985).

Rupture of the micropylar testa and the micropylar endosperm, resulting in radicle emergence, are distinct and temporally separate events during the germination of tobacco seeds (Arcila and Mohapatra, 1983; Leubner-Metzger *et al.*, 1995; Web site: <http://www.leubner.ch/>). Treatment of imbibed tobacco seeds with abscisic acid (ABA) specifically delays endosperm rupture, but not testa rupture, and results in the formation of a novel structure, consisting of the enlarging radicle covered by a sheath of greatly elongated micropylar endosperm.

As in *Cestroideae*-type seeds, β Glu also accumulates prior to endosperm rupture in *Solanoideae*-type tomato seeds (Wu *et al.*, 2000). The micropylar-covering layers of these types of seeds are organized in cap-like structures, but a distinction between testa and endosperm rupture is not possible (Watkins and Cantliffe, 1983; Watkins *et al.*, 1985; Hilhorst, 1995; Judd *et al.*, 1999; Nonogaki *et al.*, 2000). ABA also inhibits tomato endosperm rupture, i.e. radicle emergence of tomato seeds (Hilhorst, 1995; Toorop *et al.*, 2000; Wu *et al.*, 2000). Gibberellin (GA)-induced weakening of the micropylar endosperm cap is required for tomato and pepper seed germination, and the GA-deficient *gib1* mutant of tomato does not germinate without GA treatment (e.g. Watkins and Cantliffe, 1983; Watkins *et al.*, 1985; Hilhorst, 1995; Bewley, 1997a; Nonogaki *et al.*, 2000; Wu *et al.*, 2000). Weakening of the micropylar endosperm is likely to be achieved by cell-wall hydrolysis by the action of GA-induced cell-wall hydrolases.

Prior to radicle protrusion, two phases can be distinguished in tomato: (1) the early phase is not inhibited by ABA and includes ABA-insensitive endosperm weakening associated with micropylar-endosperm-specific, ABA-independent expression of endo- β -mannanase and other proteins; ABA inhibits β Glu I expression (e.g. Bewley, 1997a; Toorop *et al.*, 1998, 2000; Nonogaki *et al.*, 2000; Wu *et al.*, 2000). Endo- β -mannanase, which can hydrolyse isolated micropylar endosperm cell walls *in vitro*, appears to be necessary for endosperm weakening, but is not sufficient for the completion of tomato germination,

and its accumulation is not inhibited by ABA. (2) The late phase is critical, since it includes the final ABA-controlled step of radicle emergence. It is associated with ABA-sensitive β Glu I expression in the micropylar endosperm, and β Glu I, therefore, could contribute to radicle emergence of tomato (Wu *et al.*, 2000; Leubner-Metzger, 2003). It has been proposed that the late phase includes a second, ABA-controlled step of endosperm weakening, which thereby is a biphasic process in tomato (Toorop *et al.*, 2000).

On the molecular level, phylogenetic analyses of the multigene families of β -1,3-glucanases (β Glu) and chitinases (Chn) showed that their organization is highly conserved within the *Solanaceae* (e.g. Sperisen *et al.*, 1991; Meins *et al.*, 1992; Van Buuren *et al.*, 1992; Simmons, 1994). Based on the amino acid sequences of the mature proteins, the various β Glu and Chn have been grouped into structural classes that differ in sequence identity by at least 40–50%. At least three structural classes are identified for the highly homologous β Glu isoforms of tobacco, tomato, potato and pepper. These conserved evolutionary relationships are also manifested on the level of the hormonal regulation of the orthologous genes. For example, ethylene treatment results in the transcriptional co-induction of the class I isoforms of β Glu (β Glu I) and Chn (Chn I) in the leaves of tobacco and tomato. ABA transcriptionally down-regulates β Glu I, but not Chn I, of tobacco and tomato (Rezzonico *et al.*, 1998; Leubner-Metzger and Meins, 1999; Wu *et al.*, 2000).

Seed germination of tobacco and tomato is associated with the transcriptional induction of the β Glu I genes just prior to radicle emergence (Leubner-Metzger *et al.*, 1995; Wu *et al.*, 2000; Leubner-Metzger, 2001). This induction is highly confined to the micropylar endosperm, and β Glu I accumulation and endosperm rupture of both species are promoted by GA and inhibited by ABA. Sense transformation of tobacco with a chimeric ABA-inducible β Glu I transgene provided direct evidence that β Glu I contributes to endosperm rupture (Leubner-Metzger and Meins, 2000). In tomato, but not in tobacco, Chn I also accumulates in the micropylar endosperm prior to radicle emergence. In contrast to β Glu I, and in agreement with the situation in vegetative tissues, Chn I accumulation in imbibed tomato seeds is not inhibited by ABA. Thus, the hormonal regulation of β Glu I induction and endosperm rupture of *Cestroideae*-type (tobacco) and *Solanoideae*-type (tomato) seeds is similar, but the two seed types differ with respect to Chn expression. However, since only one type-member of each subgroup has been investigated so far, no general conclusions with respect to similarities and differences in the tissue-specific and hormonal regulation of β Glu and Chn among Solanaceous seeds can be drawn.

Materials and methods

Plant materials and germination conditions

Mature seeds of *Capsicum annuum* L. cv. Toro (ESASEM, Milano, Italy), *Physalis peruviana* L. (Wyss Samen und Pflanzen AG, Zuchwil-Solothurn, Switzerland), *Petunia hybrida* Hort. (Vilm.) (Plantania seeds, OBI and Royal Lemkes, Bleiswijk, The Netherlands), *Nicotiana sylvestris* Speg & Comes, *Nicotiana plumbaginifolia* Viv., and *Nicotiana tabacum* L. cv. Havana 425 (Agricultural Experimental Station, University of Wisconsin, Madison, Wisconsin, USA) were used for germination analyses, performed as described earlier (Leubner-Metzger *et al.*, 1998). In brief, 100–150 seeds (c. 20 seeds for *C. annuum* and *P. peruviana*) were sown in 9-cm-diameter plastic Petri dishes containing filter paper wetted with a nutrient solution (control medium) supplemented as indicated with 10 μ M *cis*-(\pm)-abscisic acid (ABA, Sigma, St. Louis, Missouri, USA) and 10 μ M gibberellin A₄ (GA₄, Sigma). Petri dishes were incubated at 24°C in continuous white light (3000 lux, Philips 'TL'D 35W/33 lamps) or in darkness. After scoring for germination, seeds were stored at -70°C for subsequent analyses.

Analysis of proteins and RNA

Procedures for extracting proteins, assays for enzyme activity, immunoblot analysis, and protein determination have been described previously (Leubner-Metzger *et al.*, 1995). In brief, β Glu and Chn activities were assayed radiometrically using reduced [³H]laminarin and [³H]chitin as the substrates, which are specifically digested by endo-type β Glu and Chn, respectively. The polyclonal rabbit anti-tobacco β Glu I antibody used for immunoblot analysis is known to detect the class I, class II and class III isoforms of *N. tabacum* and *N. sylvestris* (Neuhaus *et al.*, 1992; Beffa *et al.*, 1993; Kunz *et al.*, 2001) and is known to cross-react with the β Glu I of tomato (Wu *et al.*, 2000) and pea (Petruzzelli *et al.*, 1999). The polyclonal rabbit anti-tobacco Chn I antibody used for immunoblot analysis is known to detect the Chn I isoforms of *N. tabacum*, *N. sylvestris* (Kunz *et al.*, 2001) and tomato (Wu *et al.*, 2000). Preparation of total RNA and RNA-blot hybridization were as described by Leubner-Metzger *et al.* (1995). The probes were radiolabelled with [α -³²P]dCTP by random priming (rediprime kit; Amersham, Buckinghamshire, UK), using as templates the c. 1 kb *Pst*I fragments of the tobacco cDNAs for β Glu I (Shinshi *et al.*, 1988) and Chn I (Shinshi *et al.*, 1987), and the 1.8 kb *Eco*RI fragment of genomic DNA encoding tomato 18S ribosomal RNA (Schmidt-Puchta *et al.*, 1989). Hybridized membranes were washed at high

stringency [20 min at 62°C in 0.1% (w/v) SDS, 30 mM NaCl, 3 mM sodium citrate, pH 7.0].

Results

ABA-sensitive β Glu is induced after testa rupture, but prior to endosperm rupture in Cestroideae-type seeds

Testa rupture and endosperm rupture are distinct and temporally separate events during the germination of tobacco seeds (Arcila and Mohapatra, 1983; Leubner-Metzger *et al.*, 1995). In initial experiments we investigated whether this is also the case for other *Nicotiana* species and for *Petunia hybrida*, which all belong to the *Cestroideae* subgroup of *Solanaceae* (Judd *et al.*, 1999). Testa rupture also precedes endosperm rupture of *N. plumbaginifolia*, *N. sylvestris* and *P. hybrida* (Table 1). Seed germination of many *Nicotiana* species requires light, and, in agreement with this, neither testa rupture nor endosperm rupture was observed in *N. tabacum*, *N. plumbaginifolia* and *N. sylvestris* seeds imbibed in darkness (Table 1; Leubner-Metzger, 2003). In contrast to *N. tabacum*, *N. sylvestris*, and *P. hybrida*, light alone was not sufficient to induce testa rupture and germination of imbibed *N. plumbaginifolia* seeds, but treatment with 10 μ M GA₄ induced testa rupture and subsequent endosperm rupture (Table 1). Furthermore, as in tobacco, ABA treatment of seeds inhibited endosperm rupture, but did not affect testa rupture of the three *Nicotiana* species or of *petunia*.

Reduced [³H]laminarin, an algal β -1,3-glucan known to be specifically digested by all endo-type β Glu isoforms (Leubner-Metzger *et al.*, 1995), was utilized for the β Glu assays. As in tobacco, β Glu activity accumulated after testa rupture, but prior to endosperm rupture in germinating seeds of *N. plumbaginifolia*, *N. sylvestris* and *P. hybrida* (Table 1). No accumulation of β Glu activity was found under the conditions where these species did not germinate, i.e. in dark-imbibed *N. sylvestris* seeds and in light-imbibed *N. plumbaginifolia* seeds without GA treatment. ABA treatment not only inhibited endosperm rupture, but also inhibited the accumulation of β Glu activities in the different *Nicotiana* species and in *petunia* (Table 1; Leubner-Metzger *et al.*, 1995). β Glu activity accumulation in the micropylar endosperm during tobacco seed germination appears to be due to the transcriptional induction of the β Glu I genes. RNA-blot and immunoblot analyses presented in Fig. 1 support the view that this is also the case in the different *Nicotiana* species and in *petunia*. A tobacco β Glu I cDNA probe detected a c. 1.6 kb transcript in germinating *N.*

Table 1. Effect of gibberellin (GA) and abscisic acid (ABA) on germination and endo-type β -1,3-glucanase (β Glu) enzyme activity accumulation in imbibed Solanaceous seeds

Species	Incubation conditions	Time (hours)	Seed state ^a	Control				10 μ M GA ₄				10 μ M ABA				
				Rupture (%) ^b		β Glu	pkat ^c	Rupture (%)		β Glu	pkat	Rupture (%)		Endosperm	β Glu	pkat
				Testa	Endosperm	Testa	Endosperm	Testa	Endosperm	Testa	Endosperm	Testa	Endosperm	Testa	Endosperm	pkat
Cestroioideae subgroup of Solanaceae																
<i>Nicotiana plumbaginifolia</i>	Cont. light	25		0.0	0.0	–	–	71.2	0.0	0.01 \pm 0.00	–	–	–	–		
		45		0.0	0.0	0.01 \pm 0.00	–	95.8	23.8	0.22 \pm 0.01	–	–	0.0	0.01 \pm 0.00		
	Cont. light	45	NG				100	0.0	0.19 \pm 0.03	–	–	–	–	–		
		45	G				100	100	0.26 \pm 0.03	–	–	–	–	–		
<i>Nicotiana sylvestris</i>	Cont. light	0		0.0	0.0	0.04 \pm 0.01	–	–	–	–	–	–	–	0.04 \pm 0.01		
		60	NG	100	0.0	0.20 \pm 0.02	–	–	–	–	–	100	0.0	0.06 \pm 0.01		
	Darkness	90	G	100	100	0.79 \pm 0.05	–	–	–	–	–	100	0.0	0.10 \pm 0.01		
		90	NG	0.0	0.0	0.05 \pm 0.01	–	–	–	–	–	0.0	0.0	0.04 \pm 0.01		
<i>Petunia hybrida</i>	Cont. light	20		3.5	0.0	0.22 \pm 0.01	–	9.7	0	0.19 \pm 0.02	–	–	–	0.22 \pm 0.01		
		40		46.5	0.0	0.21 \pm 0.02	–	57.1	1.2	0.18 \pm 0.01	–	–	–	0.19 \pm 0.00		
	Cont. light	50		65.3	0.7	0.29 \pm 0.02	–	68.5	3.9	0.24 \pm 0.02	–	–	–	0.16 \pm 0.01		
		60		75.3	29.2	0.39 \pm 0.03	–	77.8	24.2	0.31 \pm 0.02	–	–	–	0.19 \pm 0.03		
	Cont. light	70		77.7	45.0	0.39 \pm 0.02	–	79.7	48.1	0.35 \pm 0.03	–	–	–	0.21 \pm 0.02		
		70	NG			0.31 \pm 0.03	–	–	–	–	–	–	–	–		
Cont. light	70	Endosperm ^d			0.45 \pm 0.01	–	–	–	–	–	–	–	–	0.20 \pm 0.01		
	70	Embryo ^d			0.37 \pm 0.04	–	–	–	–	–	–	–	–	0.10 \pm 0.00		
Solanoideae subgroup of Solanaceae																
<i>Physalis peruviana</i>	Cont. light	0			0.0	0.05 \pm 0.01	–	–	0.0	0.05 \pm 0.01	–	–	–	0.05 \pm 0.01		
		20			0.0	0.0	0.28 \pm 0.03	–	–	0.0	–	–	–	–		
	Cont. light	70			0.0	0.0	0.48 \pm 0.04	–	–	0.0	0.88 \pm 0.12	–	–	0.76 \pm 0.09		
		140			65.2	2.08 \pm 0.18	–	–	100	1.77 \pm 0.21	–	–	–	1.30 \pm 0.15		
	Cont. light	140	NG		0.0	1.23 \pm 0.11	–	–	–	–	–	–	–	–		
		140	G		100	2.53 \pm 0.27	–	–	–	–	–	–	–	–		

Cont., continuous.

^a NG, Non-germinated; G, germinated.

^b Germination of 100–150 seeds (20 seeds for *P. peruviana*) incubated at 24°C; mean values of usually 3 samples.

^c Mean values \pm SE of enzyme activities in pkat / seed (*P. peruviana*) or pkat / μ g protein (*P. hybrida*, *N. plumbaginifolia*, *N. sylvestris*).

^d β Glu activities of endosperms and embryos from control seeds (germinated) or ABA-treated seeds (non-germinated) were analysed separately.

plumbaginifolia seeds, but only upon GA treatment and not in non-germinating seeds (Fig. 1A). The rabbit anti-tobacco β Glu I antibody used for immunoblot analysis is known to detect the class I, class II and class III β Glu isoforms of *N. tabacum* and *N. sylvestris* (Neuhaus *et al.*, 1992; Beffa *et al.*, 1993). This antibody detected the accumulation of immunoreactive bands in germinating seeds of *N. sylvestris* imbibed in the light (c. 33 kDa; data not shown) and of *N. plumbaginifolia* imbibed in the light and treated with GA (c. 34 kDa; Fig. 1B). The sizes of these bands are in agreement with the known sizes of the β Glu I of these species (Castresana *et al.*, 1990; Gheysen *et al.*, 1990; Neuhaus *et al.*, 1992), and the bands were not detected in seeds imbibed under conditions that prevent or inhibit germination. The initial β Glu activities in seeds of the three *Nicotiana* species are essentially background level (Table 1), and the β Glu I accumulation during imbibition is localized exclusively in the endosperm (data not shown; Leubner-Metzger *et al.*, 1995). In contrast to *Nicotiana* seeds, the initial β Glu activities of petunia seeds were considerably higher, and β Glu activity accumulation during imbibition was localized in the endosperm and in the embryo (Table 1). Figure 1C shows that the anti-tobacco β Glu I antibody detects several immunoreactive bands (27, 30, 35, 43, 47 and 50 kDa in size) in the protein extracts of germinating petunia seeds. The 27, 30, 47 and 50 kDa bands did not show appreciable regulation with respect to the germination process and to the hormone treatments. In contrast, the 35 kDa and the 43 kDa bands appeared to be down-regulated by ABA and up-regulated by imbibition and/or GA, in agreement with the β Glu activities. These two putative β Glu isoforms are located in the endosperm and in the embryo, with a predominance of the 43 kDa bands in the embryo. Taken together, these findings in four species suggested that the accumulation of ABA-sensitive β Glu I in the endosperm after testa rupture, but prior to endosperm rupture, appears to be a common phenomenon during the germination of *Cestroideae*-type seeds.

β Glu and Chn are co-induced in the roots of Solanaceous seedlings as a post-germination event

β Glu I induction in germinating tobacco seeds prior to endosperm rupture is confined to the micropylar endosperm and is absent from the lateral endosperm and the embryo (Fig. 2A; Leubner-Metzger *et al.*, 1995). When we extended our studies to germinated tobacco seedlings at 90 h and 150 h, we found an additional, post-germination induction phase of β Glu I that is localized in the root, but not appreciably in the shoot of the seedlings (Fig. 2A; Table 2). Chn is not induced prior to the completion of endosperm

rupture of *N. tabacum* (Fig. 2B, C; Leubner-Metzger *et al.*, 1995) and *N. sylvestris* (data not shown). However, in germinated tobacco seedlings at 90 h and 150 h, the mRNA and the 32 kDa and 34 kDa antigens of Chn I are detected (Fig. 2B, C). As for the β Glu I, Chn I accumulation is also mainly localized to the root of the tobacco seedlings (Table 2). In contrast to β Glu I, which is transcriptionally down-regulated by ABA in seeds and in vegetative tissues, Chn I expression in vegetative tissues is not down-regulated by ABA (Leubner-Metzger *et al.*, 1995; Rezzonico *et al.*, 1998). This was also the case in ABA-treated tobacco seeds and, since ABA delays endosperm rupture, this leads to the finding that Chn I accumulates prior to endosperm rupture in ABA-treated seeds (Fig. 2B, C). This is in contrast to seeds imbibed in medium without ABA added, and the Chn I levels found in 150 h seedlings were similar to those found in 150 h seeds treated with ABA (Fig. 2B). In contrast to tobacco, Chn I is also induced prior to germination in tomato seeds in medium lacking ABA (Wu *et al.*, 2000). As in tobacco, β Glu and Chn activity accumulations are associated with the root, but appreciably less with the shoot, of tomato and pepper seedlings (Table 2). Thus, a post-germination co-induction of β Glu and Chn in the root of seedlings seems to be a general phenomenon in both subgroups of the *Solanaceae*.

Temporal and spatial pattern of β Glu and Chn accumulation are distinct among Solanoideae-type seeds

Tomato, *Physalis peruviana* and *Capsicum annuum* belong to the *Solanoideae* subgroup of *Solanaceae* and are characterized by flattened, discoid seeds with curved embryos, and the micropylar endosperm and testa layers of pepper and tomato form a cap-like structure (Watkins and Cantliffe, 1983; Watkins *et al.*, 1985; Judd *et al.*, 1999). β Glu I and Chn I are both co-expressed in the micropylar endosperm of tomato prior to radicle emergence (Wu *et al.*, 2000). Using highly sensitive radiometric assays with [3 H]laminarin and [3 H]chitin as substrates, respectively, similar contents and kinetics of specific β Glu and Chn activities were obtained in tomato seeds. Figure 3 shows that by using the same highly sensitive radiometric assays, only the accumulation of β Glu activity, but not of Chn activity, was detected in germinating pepper seeds. The onset of radicle protrusion of imbibed pepper seed populations was not detected in the 72 h samples, but started slightly later. As in tomato, β Glu activity accumulates in the micropylar cap of pepper seeds prior to its rupture by the protruding radicle and germination is inhibited by ABA (Fig. 3B). In contrast to tomato, where β Glu accumulation is confined to the micropylar




Figure 1. Hormone-regulated accumulation of tobacco β Glu I-related mRNAs and antigens during the germination of *Cestroideae*-type *Solanaceae* seeds. (A) RNA-blot hybridization of total RNA (25 μ g/lane) prepared from *Nicotiana plumbaginifolia* seeds imbibed in continuous light either in the absence (Control) or presence of 10 μ M GA₄ (GA); only GA-treated seeds germinate. The RNA-blot was hybridized with a cDNA probe for tobacco β Glu I. (B) Immunoblot analysis of *N. plumbaginifolia* seed extracts (80 μ g protein/lane) probed with the rabbit anti-tobacco β Glu I antibody. G = germinated seeds only; NG = ungerminated seeds only. β Glu I = 10 ng of the authentic 33 kDa tobacco enzyme. No signals were obtained in control blots with rabbit anti-tobacco Chn I antibody or with rabbit pre-immune serum. (C) Immunoblot analysis of *Petunia hybrida* seed extracts using the anti-tobacco β Glu I antibody; GA = 10 μ M GA₄; ABA = 10 μ M ABA, End = endosperm, Emb = embryo; analysis as described in (B) with the difference that 50 μ g protein/lane (entire seeds) and 30 μ g protein/lane (End, Emb) were applied.




Figure 2. Tissue specificity and regulation by abscisic acid (ABA) of βGlu I and Chn I in imbibed seeds and in seedlings of *Nicotiana tabacum*. (A) RNA-blot hybridization of total RNA (25 μg/lane) prepared from seeds prior to endosperm rupture (55 h) during imbibition in continuous light and from 90 h seedlings. The RNA-blot was hybridized with a cDNA probe for tobacco βGlu I, and a probe for 18S rRNA was used as a loading standard. Em = embryo; ME = micropylar endosperm; LE = lateral endosperm; S = shoot; R = root; E = 90 h endosperm remnant. (B) Immunoblot analysis of seed and seedling extracts (80 μg protein/lane) probed with the rabbit anti-tobacco Chn I antibody. Seeds and seedlings were imbibed in continuous light either in the absence (Control) or presence of 10 μM ABA. CHN I = 10 ng of the authentic 32 kDa and 34 kDa tobacco enzymes. (C) RNA-blot hybridization of total RNA prepared from seeds and seedlings incubated in medium without (Con) or with 10 μM ABA added (ABA). The RNA-blot was hybridized with probes for tobacco Chn I and 18S rRNA as described in (A).

Table 2. Tissue-specificity of β-1,3-glucanase (βGlu) and chitinase (Chn) enzyme activity in Solanaceous seedlings

Species ^b	βGlu ^a			Chn ^a		
	Root (R)	Shoot (S)	R/S ratio ^c	Root (R)	Shoot (S)	R/S ratio
Pepper	3.3 ± 0.6	0.1 ± 0.0	33	8.2 ± 0.9	0.1 ± 0.0	82
Tomato	1.0 ± 0.5	0.1 ± 0.0	10	–	–	–
Tobacco	11.7 ± 1.1	2.0 ± 0.6	6	31.1 ± 5.5	6.8 ± 2.6	5

^a Mean values ± SE of enzyme activities in pkat/μg protein of four protein samples, each from 10–15 seedling tissues.

^b Seedlings from germinated seeds incubated for 150 h (tobacco) or 160 h (pepper, tomato) in continuous light at 24°C.

^c Ratio between the specific enzyme activities of the seedling root and shoot tissues.

endosperm, βGlu activity also accumulated in the embryo (Fig. 3A) and the rest of the pepper seed (Fig. 3C). The βGlu activity accumulation in all three compartments was inhibited by ABA (Fig. 3). βGlu activity also accumulated in imbibed seeds of *P. peruviana* and the enzyme activity accumulation was only partially inhibited by ABA (Table 1).

To further investigate the tissue- and species-specific accumulation of βGlu isoforms, we performed immunoblot analyses with the different pepper seed tissues. The rabbit anti-tobacco βGlu I antibody, which is known to also detect the 35 kDa tomato βGlu I in the micropylar endosperm (Wu *et al.*, 2000), detected a distinct temporal and spatial pattern




Figure 3. Tissue specificity and regulation by ABA of β Glu and Chn activities during the seed germination of *Capsicum annuum*, measured in (A) embryo, (B) micropylar cap and (C) the rest of the seed. Populations of 20 seeds were imbibed in continuous light, either in the absence (Control) or presence of 100 μ M ABA (ABA), and seeds were dissected at the times indicated. Specific β Glu and Chn activities are expressed in pkat/ μ g protein. Mean values \pm SE are presented from four to ten replicate samples at each time point. Radicle protrusion of imbibed pepper seeds: 0% until 72 h, 11.6% at 96 h, 100% at 120 h (Control) and 0% for ABA-treated seeds at 120 h. At 96 h non-germinated (N) and germinated (G) seeds were measured separately.

of immunoreactive bands in the different seed compartments and in seedlings (Fig. 4). A major, *c.* 39.5 kDa band increased in intensity in the embryo prior to and during endosperm rupture, but was not detected in the micropylar cap prior to endosperm rupture or in the rest of the seed. The 39.5 kDa band was also present in the micropylar cap of 120 h germinated seeds, but not in ABA-treated ungerminated seeds. The appearance of the 39.5 kDa band in the embryo was inhibited by ABA (Fig. 4A), which also inhibited β Glu activity accumulation (Fig. 3A). The 39.5 kDa band was the only band detected in seedling shoots and was absent from seedling roots (Fig. 4A). Additional bands, *c.* 27 kDa and 31 kDa in size, were detected at roughly constitutive levels in all seed tissues (Fig. 4). The 27 kDa band was especially strong in the embryo, but was absent from seedlings. Roughly constitutive levels of a *c.* 24 kDa band were found in the endosperm, but not in embryos or seedlings. Treatment with ABA did not affect the levels of the 24, 27 and 31 kDa bands, but the transition from the embryo state to the seedling state was associated with the disappearance of these bands. A *c.* 33 kDa band was detected in seedling roots. Thus, a complex tissue-specific pattern of putative β Glu was detected in imbibed seeds and in seedlings of pepper. In contrast to pepper, the rabbit anti-tobacco β Glu I antibody did not detect any antigen in protein extracts from seed extracts of *P. peruviana*. Finally, in agreement with the finding that Chn activity did not accumulate during pepper seed germination, the rabbit anti-tobacco Chn I antibody did not detect immunoreactive bands in imbibed pepper seeds (data not shown); but it did detect Chn I in tomato seeds (Wu *et al.*, 2000).

Discussion

Two developmentally regulated sites of β Glu induction seem to be common among Solanaceous species: (1) β Glu accumulation in imbibed seeds prior to endosperm rupture, and (2) β Glu accumulation in roots of young seedlings as a post-germination event. Whereas co-induction of Glu and Chn in seedling roots appears to be a common phenomenon, significant differences exist among Solanaceous species during seed germination. The various β Glu isoforms of tobacco, tomato, potato, pepper and other species have been classified into at least three structural classes that differ by a minimum of 40–50% in amino acid sequence identity of the mature proteins (Meins *et al.*, 1992; Simmons, 1994; Leubner-Metzger and Meins, 1999). The endo-type class I enzymes (β Glu I) include the highly homologous 33 kDa isoforms of *N. tabacum* and *N. sylvestris* (Sperisen *et al.*, 1991; Neuhaus *et al.*, 1992), the 34 kDa

isoforms of *N. plumbaginifolia* (*Gn2* and *Gn1* share 98% and 76% amino acid sequence identity with tobacco β Glu I, respectively; Castresana *et al.*, 1990; Gheysen *et al.*, 1990), the 35 kDa tomato isoform Glu B (*c.* 90% amino acid sequence identity with tobacco β Glu I; Van Kan *et al.*, 1992; Wu *et al.*, 2000), and the β Glu I of pepper exhibits a similarly high degree of sequence identity (Kim and Hwang, 1997; Jung and Hwang, 2000b; Pflieger *et al.*, 2001). These β Glu I isoforms are usually basic, intracellular proteins, but there is strong evidence that they can also be secreted (Simmons, 1994; Leubner-Metzger and Meins, 1999). In contrast to β Glu I, the class II and III β Glu are secreted into the apoplast. Characteristic differences between class I and II β Glu genes are also evident at the level of gene regulation by ethylene, ABA, salicylic acid, pathogens and other factors. β Glu I gene expression is induced transcriptionally in vegetative tissues and in seeds by ethylene, and down-regulated by ABA (e.g. Leubner-Metzger *et al.*, 1995; Rezzonico *et al.*, 1998; Jung and Hwang, 2000b; Wu *et al.*, 2000). As for β Glu, similar structural classes exist for the Chn isoforms, and β Glu I and Chn I are often co-induced, e.g. in leaves by ethylene or pathogens. In contrast to β Glu I, Chn I is not transcriptionally down-regulated by ABA and does not accumulate in the micropylar endosperm of germinating tobacco seeds (Leubner-Metzger *et al.*, 1995), but does accumulate in the micropylar endosperm of germinating tomato seeds (Wu *et al.*, 2000). The present study shows that *Cestroideae*- and *Solanoideae*-type seeds not only differ in their morphology (Judd *et al.*, 1999), but that they also exhibit common and distinct aspects with regard to the hormonal and developmental regulation of β Glu and Chn.

β Glu induction in the endosperm, and endosperm rupture but not testa rupture, are inhibited by ABA in Cestroideae-type seeds

Our most intriguing findings with *Cestroideae*-type seeds (three *Nicotiana* species and petunia) are the common distinction between testa rupture and endosperm rupture, and that ABA-sensitive β Glu accumulation occurs prior to endosperm rupture, but after testa rupture. As in tobacco (Arcila and Mohapatra, 1983; Leubner-Metzger *et al.*, 1995), testa rupture and endosperm rupture of *N. sylvestris*, *N. plumbaginifolia* and petunia are distinct and sequential events. As in tobacco, ABA does not appreciably inhibit testa rupture, but inhibits β Glu accumulation in the endosperm and endosperm rupture (Table 3). GA treatment is required to release dormancy and induce β Glu accumulation and germination of *N. plumbaginifolia* seeds imbibed in the light and *N. sylvestris* seeds imbibed in darkness. These results are




Figure 4. Tissue specificity and regulation by ABA of β Glu antigens during germination of *Capsicum annuum* seeds, in (A) embryo and 160 h-seedling, (B) micropylar cap and (C) the rest of the seed. Immunoblot analysis of seed and seedling extracts (40 μ g protein/lane) probed with the rabbit anti-tobacco β Glu I antibody, which recognizes all known tobacco β -1,3-glucanases and is also known to cross-react with tomato β Glu I (Neuhaus *et al.*, 1992; Beffa *et al.*, 1993; Petruzzelli *et al.*, 1999; Wu *et al.*, 2000; Kunz *et al.*, 2001). No signals were obtained in control blots with rabbit anti-tobacco Chn I antibody or with rabbit pre-immune serum. Extracts and germination are as described in Fig. 3. R = root, S = shoot.

in agreement with the roles of ABA and GA in regulating dormancy and germination of *Nicotiana* species (e.g. Leubner-Metzger *et al.*, 1996; Frey *et al.*, 1999; Grappin *et al.*, 2000; Leubner-Metzger, 2001) and of petunia (Sink, 1984; Girard, 1990). The highly homologous 34 kDa and 33 kDa β Glu I isoforms of *N. plumbaginifolia* (Castresana *et al.*, 1990; Gheysen *et al.*, 1990) and *N. sylvestris* (Sperisen *et al.*, 1991; Neuhaus *et al.*, 1992), respectively, accumulate after testa rupture, but prior to endosperm rupture. As in tobacco, germination and β Glu I accumulation seem to depend on the light/GA pathway and are inhibited by ABA (Leubner-Metzger *et al.*, 1996; Leubner-Metzger, 2001). No sequence information is available for petunia β Glu isoforms, but the anti-tobacco β Glu I antibody detected several constitutively expressed immunoreactive bands, as well as two immunoreactive bands, 35 kDa and 43 kDa in size, that exhibit ABA-inhibited regulation. In contrast to *Nicotiana* seeds, enzyme activity and antigen accumulation in petunia are present in both endosperm and embryo. It seems likely, but not

proven, that the 35 kDa band that accumulates in association with endosperm rupture is a petunia β Glu I. Taken together, these findings suggest that the accumulation of β Glu I prior to endosperm rupture and the regulation of β Glu I expression by light/GA and ABA could be an evolutionarily conserved developmental event in *Cestroidae*-type seeds (Leubner-Metzger, 2003).

Distinct expression pattern of β Glu and Chn isoforms in germinating Solanaceous seeds

ABA also inhibits endosperm rupture of *Solanoideae*-type seeds such as tomato (Hilhorst, 1995; Toorop *et al.*, 2000; Wu *et al.*, 2000) and pepper, but not *P. peruviana* (this study). As in *Cestroidae*-type seeds, β Glu also accumulates prior to endosperm rupture of tomato, pepper and *Physalis* seeds (Table 3). As in tobacco, β Glu accumulation in the micropylar endosperm of tomato and pepper is inhibited by ABA. In contrast to tobacco and tomato, β Glu accumulation in pepper and petunia seeds is not

Table 3. β-1,3-Glucanase (βGlu) and chitinase (Chn) activity induction in imbibed Solanaceous seeds and in young seedlings

Species ^a	Seeds prior to endosperm rupture						Seedling roots	
	βGlu induction		Chn induction		Inhibition by ABA of endosperm		βGlu and Chn co-induction	
	Endosperm ^b	Embryo	Endosperm	Embryo	βGlu induction	rupture		
<i>Cestroideae</i> subgroup of <i>Solanaceae</i>								
<i>Nicotiana tabacum</i>	+ ^c	-	-	-	+	+	+	+
<i>Nicotiana sylvestris</i>	+	-	-	-	+	+	+	+
<i>Nicotiana plumbaginifolia</i>	+	-	n.d.	n.d.	+	+	+	n.d.
<i>Petunia hybrida</i>	+	+	n.d.	n.d.	+	+	+	n.d.
<i>Solanoidae</i> subgroup of <i>Solanaceae</i>								
<i>Lycopersicon esculentum</i>	+	-	+	+	+	+	+	+
<i>Capsicum annuum</i>	+	+	-	-	+	+	+	+
<i>Physalis peruviana</i>	+	n.d. ^d	n.d.	n.d.	(-) ^e	+	+	n.d.

^a Based on this study and on additional results for *Nicotiana* species (Vögel-Lange *et al.*, 1994; Leubner-Metzger *et al.*, 1995; Kunz *et al.*, 2001) and tomato (Wu *et al.*, 2000).

^b Localization of βGlu induction in the micropylar part of the endosperm has been demonstrated for *N. tabacum*, *N. sylvestris*, *L. esculentum*, and *C. annuum*; for the other species only accumulation in the overall endosperm has been demonstrated.

^c A '+' sign means presence and a '-' sign absence of a feature or process.

^d n.d. = not determined.

^e No inhibition by ABA of overall seed βGlu activity accumulation, which does not *per se* exclude the existence of an ABA-responsive βGlu isoform in the endosperm.

confined exclusively to the micropylar tissue, but also accumulates in other seed parts, especially in the embryo. At least ten basic and acidic β Glu isoforms have been described in healthy and infected vegetative pepper tissues, corresponding to sizes of 25, 27, 29, 34, and 36 kDa in denaturing gels (e.g. Kim and Hwang, 1994, 1997; Egea *et al.*, 1999). The limited sequence information on pepper available in databases includes a 39.3 kDa ethylene-inducible β Glu I (Jung and Hwang, 2000a, b; Pflieger *et al.*, 2001). Using the anti-tobacco β Glu I antibody, we detected a major *c.* 39.5 kDa band in the embryo prior to and during germination, and in seedling shoots. This band is not present during early imbibition, is not induced in the micropylar cap prior to endosperm rupture, nor in the rest of the seed or in seedling roots. Its appearance is inhibited by ABA and it corresponds in size to the ethylene-inducible 39.3 kDa β Glu I described by Jung and Hwang (2000b). The anti-tobacco β Glu I antibody also detected several other immunoreactive bands that were 24, 27 and 31 kDa in size, roughly constitutive and not affected by ABA. The nature of these bands is not known, and the limited sequence information, together with expected cultivar differences, does not allow an unambiguous identification of these isoforms. ABA-sensitive β Glu activity accumulation was especially strong in the micropylar endosperm of pepper seeds, but none of the detected bands correlated with this activity pattern. Thus, a serologically distinct ABA-sensitive β Glu isoform must account for the β Glu accumulation in the micropylar cap of imbibed pepper seeds. The anti-tobacco β Glu I antibody also detected a *c.* 33 kDa band, but only in seedling roots. Considering the known serological relatedness of the tobacco and tomato β Glu I with the 34 kDa pepper β Glu I (e.g. Kim and Hwang, 1997; Wu *et al.*, 2000), we propose that this is a pepper β Glu I, probably the basic 34 kDa isoform described by Kim and Hwang (1997). The tissue-specific and ABA-regulated differences in band patterns suggest that several distinct β Glu isoforms account for the β Glu activities of pepper and petunia. ABA did not inhibit overall β Glu activity accumulation in seeds of *P. peruviana*; however, it is still a reasonable, but unproven, hypothesis that several distinct β Glus accumulate and that, among them, are ABA-sensitive and -insensitive isoforms. The finding that the anti-tobacco β Glu I antibody did not detect any signals in seed extracts of *P. peruviana* is a complication for testing this hypothesis. Prior to endosperm rupture, β Glu I expression in *Nicotiana* and tomato seeds is confined to the micropylar endosperm, whereas in pepper and petunia seeds, distinct isoforms confer β Glu activity accumulation in several tissues (this study; Leubner-Metzger *et al.*, 1995; Wu *et al.*, 2000).

Chn is not expressed in *Nicotiana* and pepper seeds prior to endosperm rupture, but Chn I is expressed in the micropylar endosperm of tomato (Table 3). Interestingly, jasmonates or wounding induce Chn I expression in the micropylar endosperm of tomato (Wu and Bradford, 2002), but do not induce β Glu I in tomato (Wu and Bradford, 2002) or tobacco seeds (Leubner-Metzger, personal communication). Thus, β Glu I and Chn I induction in the micropylar endosperm may be regulated by distinct signalling pathways. Therefore, we propose that signalling of ABA-sensitive β Glu induction in the micropylar endosperm is an evolutionarily conserved pathway that is widespread in Solanaceous seeds and serves a developmental function during endosperm rupture. This is in agreement with direct evidence obtained by sense transformation of tobacco, which showed that β Glu I contributes to endosperm rupture (Leubner-Metzger and Meins, 2000). In contrast, β Glu expression in other seed tissues and Chn expression in seeds occur only in some Solanaceous species, is regulated by distinct and diverse pathways, and therefore might serve different developmental or pathogenesis-related functions (Wu and Bradford, 2002; Leubner-Metzger, 2003).

Evolutionarily conserved induction of β Glu in seedling roots of endospermic and non-endospermic species

We found that a post-germination induction of β Glu and Chn occurred mainly in the roots of young seedlings and appears to be an additional evolutionarily conserved site of β Glu I induction (Table 3). In contrast to the micropylar endosperm, β Glu I and Chn I are co-induced in the roots of young tobacco seedlings. β Glu and Chn are also co-induced in the roots of tomato and pepper seedlings, and appear to be the class I isoforms. β Glu I is also induced in seedlings of *N. sylvestris*, and increased β Glu I and Chn I levels are present in older seedlings of this species (Vögeli-Lange *et al.*, 1994; Kunz *et al.*, 1996, 2001). β Glu I and Chn I accumulate at high concentrations in the roots and in lower leaves of mature, healthy Solanaceous plants (Van de Rhee *et al.*, 1993; Beerhues and Kombrink, 1994; Vögeli-Lange *et al.*, 1994; Leubner-Metzger and Meins, 1999). The establishment of this gradient of β Glu I and Chn I expression is regulated by ethylene. In agreement with these findings, endogenous ethylene also induces the accumulation of a β Glu I in the root, but not the shoot, of pea seedlings directly after germination (Petruzzelli *et al.*, 1999, 2000, 2003). Within the pea root, ethylene responsiveness and β Glu I accumulation are localized to the elongation and differentiation zones where the root hairs form. In contrast to Solanaceous seeds, pea seeds are non-

dormant and non-endospermic. This supports the view that the post-germination induction of β Glu I (and possibly also Chn I) is widespread among dicot seedlings and represents an evolutionarily conserved event (Leubner-Metzger, 2003).

Acknowledgements

We thank Waltraud Schmidt-Puchta for kindly providing plasmids and Sjoerd van Eeden (Friedrich Miescher Institute, Basel, Switzerland) for care of plants. This research is supported by a grant from the Deutsche Forschungsgemeinschaft (LE 720/3) to G.L.-M., which is gratefully acknowledged.

References

- Arcila, J. and Mohapatra, S.C. (1983) Development of tobacco seedling. 2. Morphogenesis during radicle protrusion. *Tobacco Science* **27**, 35–40.
- Beerhues, L. and Kombrink, E. (1994) Primary structure and expression of mRNAs encoding basic chitinase and 1,3- β -glucanase in potato. *Plant Molecular Biology* **24**, 353–367.
- Beffa, R.S., Neuhaus, J.-M. and Meins, F. (1993) Physiological compensation in antisense transformants: Specific induction of an 'ersatz' glucan endo-1,3- β -glucosidase in plants infected with necrotizing viruses. *Proceedings of the National Academy of Sciences, USA* **90**, 8792–8796.
- Bewley, J.D. (1997a) Breaking down the walls – a role for endo- β -mannanase in release from seed dormancy? *Trends in Plant Science* **2**, 464–469.
- Bewley, J.D. (1997b) Seed germination and dormancy. *Plant Cell* **9**, 1055–1066.
- Bihlmeier, M. (1927) Der Einfluss der Vorquellung und der Samenschale auf die Keimung lichtgeförderter Samen. *Jahrbücher für Wissenschaftliche Botanik* **67**, 702–732.
- Castresana, C., De Carvalho, F., Gheysen, G., Habets, M., Inze, D. and van Montagu, M. (1990) Tissue-specific and pathogen-induced regulation of a *Nicotiana plumbaginifolia* β -1,3-glucanase gene. *Plant Cell* **2**, 1131–1143.
- Egea, C., Dickinson, M.J., Candela, M. and Candela, E.M. (1999) β -1,3-Glucanase isoenzymes and genes in resistant and susceptible pepper (*Capsicum annuum*) cultivars infected with *Phytophthora capsici*. *Physiologia Plantarum* **107**, 312–318.
- Frey, A., Audran, C., Marin, E., Sotta, B. and Marion-Poll, A. (1999) Engineering seed dormancy by the modification of zeaxanthin epoxidase gene expression. *Plant Molecular Biology* **39**, 1267–1274.
- Gheysen, G., Inze, D., Soetaert, P., van Montagu, M. and Castresana, C. (1990) Sequence of a *Nicotiana plumbaginifolia* β -1,3-glucanase gene encoding a vacuolar isoform. *Nucleic Acids Research* **18**, 6685.
- Girard, J. (1990) Study of the inheritance of seed primary dormancy and the ability to enter secondary dormancy in *Petunia*: influence of temperature, light and gibberellic acid on dormancy. *Plant Cell and Environment* **13**, 827–832.
- Grappin, P., Bouinot, D., Sotta, B., Miginiac, E. and Jullien, M. (2000) Control of seed dormancy in *Nicotiana plumbaginifolia*: post-imbibition abscisic acid synthesis imposes dormancy maintenance. *Planta* **210**, 279–285.
- Hilhorst, H.W.M. (1995) A critical update on seed dormancy. I. Primary dormancy. *Seed Science Research* **5**, 61–73.
- Hilhorst, H.W.M. and Downie, B. (1996) Primary dormancy in tomato (*Lycopersicon esculentum* cv. MoneyMaker): Studies with the *sitiens* mutant. *Journal of Experimental Botany* **47**, 89–97.
- Judd, W.S., Campbell, C.S., Kellog, E.A. and Stevens, P.F. (1999) *Plant systematics: a phylogenetic approach*. Sunderland, Massachusetts, USA, Sinauer Associates, Inc.
- Jung, H.W. and Hwang, B.K. (2000a) Isolation, partial sequencing, and expression of pathogenesis-related cDNA genes from pepper leaves infected by *Xanthomonas campestris* pv. *vesicatoria*. *Molecular Plant-Microbe Interactions* **13**, 136–142.
- Jung, H.W. and Hwang, B.K. (2000b) Pepper gene encoding a basic β -1,3-glucanase is differentially expressed in pepper tissues upon pathogen infection and ethephon or methyl jasmonate treatment. *Plant Science* **159**, 97–106.
- Kim, Y.J. and Hwang, B.K. (1994) Differential accumulation of β -1,3-glucanase and chitinase isoforms in pepper stems infected by compatible and incompatible isolates of *Phytophthora capsici*. *Physiological and Molecular Plant Pathology* **45**, 195–209.
- Kim, Y.J. and Hwang, B.K. (1997) Isolation of a basic 34 kilodalton β -1,3-glucanase with inhibitory activity against *Phytophthora capsici* from pepper stems. *Physiological and Molecular Plant Pathology* **50**, 103–115.
- Kincaid, R.R. (1935) The effects of certain environmental factors on the germination of Florida cigar-wrapper tobacco seeds. *Technical Bulletin of the University of Florida Agricultural Experimental Station* **277**, 5–47.
- Koornneef, M., Bentsink, L. and Hilhorst, H. (2002) Seed dormancy and germination. *Current Opinion in Plant Biology* **5**, 33–36.
- Kunz, C., Schöb, H., Stam, M., Kooter, J.M. and Meins, F. (1996) Developmentally regulated silencing and reactivation of tobacco chitinase transgene expression. *Plant Journal* **10**, 437–450.
- Kunz, C., Schöb, H., Leubner-Metzger, G., Glazov, E. and Meins, F. (2001) β -1,3-Glucanase and chitinase transgenes in hybrids show distinctive and independent patterns of posttranscriptional gene silencing. *Planta* **212**, 243–249.
- Leubner-Metzger, G. (2001) Brassinosteroids and gibberellins promote tobacco seed germination by distinct pathways. *Planta* **213**, 758–763.
- Leubner-Metzger, G. (2002) Seed after-ripening and over-expression of class I β -1,3-glucanase confer maternal effects on tobacco testa rupture and dormancy release. *Planta* **215**, 959–968.
- Leubner-Metzger, G. (2003) Functions of β -1,3-glucanases during seed germination, dormancy release and after-ripening. *Seed Science Research* **13**, 17–34.
- Leubner-Metzger, G. and Meins, F. (1999) Functions and regulation of plant β -1,3-glucanases (PR-2). pp. 49–76 in

- Datta, S.K.; Muthukrishnan, S. (Eds) *Pathogenesis-related proteins in plants*. Boca Raton, Florida, CRC Press.
- Leubner-Metzger, G. and Meins, F.** (2000) Sense transformation reveals a novel role for class I β -1,3-glucanase in tobacco seed germination. *Plant Journal* **23**, 215–221.
- Leubner-Metzger, G., Fründt, C., Vögeli-Lange, R. and Meins, F.** (1995) Class I β -1,3-glucanase in the endosperm of tobacco during germination. *Plant Physiology* **109**, 751–759.
- Leubner-Metzger, G., Fründt, C. and Meins, F.** (1996) Effects of gibberellins, darkness and osmotica on endosperm rupture and class I β -1,3-glucanase induction in tobacco seed germination. *Planta* **199**, 282–288.
- Leubner-Metzger, G., Petruzzelli, L., Waldvogel, R., Vögeli-Lange, R. and Meins, F.** (1998) Ethylene-responsive element binding protein (EREBP) expression and the transcriptional regulation of class I β -1,3-glucanase during tobacco seed germination. *Plant Molecular Biology* **38**, 785–795.
- Liptay, A. and Schopfer, P.** (1983) Effect of water stress, seed coat restraint, and abscisic acid upon different germination capabilities of two tomato lines at low temperature. *Plant Physiology* **73**, 935–938.
- Meins, F., Neuhaus, J.-M., Sperisen, C. and Ryals, J.** (1992) The primary structure of plant pathogenesis-related glucanohydrolases and their genes. pp. 245–282 in Boller, T.; Meins, F. (Eds) *Genes involved in plant defense*. Vienna, Springer-Verlag.
- Neuhaus, J.M., Flores, S., Keefe, D., Ahl-Goy, P. and Meins, F.** (1992) The function of vacuolar β -1,3-glucanase investigated by antisense transformation. Susceptibility of transgenic *Nicotiana sylvestris* plants to *Cercospora nicotianae* infection. *Plant Molecular Biology* **19**, 803–813.
- Nonogaki, H., Gee, O.H. and Bradford, K.J.** (2000) A germination-specific endo- β -mannanase gene is expressed in the micropylar endosperm cap of tomato seeds. *Plant Physiology* **123**, 1235–1245.
- Petruzzelli, L., Kunz, C., Waldvogel, R., Meins, F. and Leubner-Metzger, G.** (1999) Distinct ethylene- and tissue-specific regulation of β -1,3-glucanases and chitinases during pea seed germination. *Planta* **209**, 195–201.
- Petruzzelli, L., Coraggio, I. and Leubner-Metzger, G.** (2000) Ethylene promotes ethylene biosynthesis during pea seed germination by positive feedback regulation of 1-aminocyclopropane-1-carboxylic acid oxidase. *Planta* **211**, 144–149.
- Petruzzelli, L., Sturaro, M., Mainieri, D. and Leubner-Metzger, G.** (2003) Calcium requirement for ethylene-dependent responses involving 1-aminocyclopropane-1-carboxylic acid oxidase in radicle tissues of germinated pea seeds. *Plant Cell and Environment* (in press).
- Pflieger, S., Palloix, A., Caranta, C., Blattes, A. and Lefebvre, V.** (2001) Defense response genes co-localize with quantitative disease resistance loci in pepper. *Theoretical and Applied Genetics* **103**, 920–929.
- Rezzonico, E., Flury, N., Meins, F. and Beffa, R.** (1998) Transcriptional down-regulation by abscisic acid of pathogenesis-related β -1,3-glucanase genes in tobacco cell cultures. *Plant Physiology* **117**, 585–592.
- Schmidt-Puchta, W., Kütemeier, G., Günther, I., Haas, B. and Sängler, H.L.** (1989) Cloning and sequence analysis of the 18S ribosomal RNA gene of tomato and a secondary structure model of the 18S rRNA of angiosperms. *Molecular and General Genetics* **219**, 17–25.
- Shinshi, H., Mohnen, D. and Meins, F.** (1987) Regulation of a plant pathogenesis-related enzyme inhibition of chitinase and chitinase mRNA accumulation in cultured tobacco tissues by auxin and cytokinin. *Proceedings of the National Academy of Sciences, USA* **84**, 89–93.
- Shinshi, H., Wenzler, H., Neuhaus, J.-M., Felix, G., Hofsteenge, J. and Meins, F.** (1988) Evidence of N- and C-terminal processing of a plant defense-related enzyme: Primary structure of tobacco prepro- β -1,3-glucanase. *Proceedings of the National Academy of Sciences, USA* **85**, 5541–5545.
- Simmons, C.R.** (1994) The physiology and molecular biology of plant 1,3- β -D-glucanases and 1,3;1,4- β -D-glucanases. *Critical Reviews in Plant Sciences* **13**, 325–387.
- Sink, K.C.** (1984) *Petunia*. Berlin, Springer-Verlag.
- Sperisen, C., Ryals, J. and Meins, F.** (1991) Comparison of cloned genes provides evidence for intergenomic exchange of DNA in the evolution of a tobacco glucan endo-1,3- β -glucosidase gene family. *Proceedings of the National Academy of Sciences, USA* **88**, 1820–1824.
- Toorop, P.E., van Aelst, A.C. and Hilhorst, H.W.M.** (1998) Endosperm cap weakening and endo- β -mannanase activity during priming of tomato (*Lycopersicon esculentum* cv. Moneymaker) seeds are initiated upon crossing a threshold water potential. *Seed Science Research* **8**, 483–491.
- Toorop, P.E., van Aelst, A.C. and Hilhorst, H.W.M.** (2000) The second step of the biphasic endosperm cap weakening that mediates tomato (*Lycopersicon esculentum*) seed germination is under control of ABA. *Journal of Experimental Botany* **51**, 1371–1379.
- Van Buuren, M., Neuhaus, J.M., Shinshi, H., Ryals, J. and Meins, F.** (1992) The structure and regulation of homologous tobacco endochitinase genes of *Nicotiana sylvestris* and *N. tomentosiformis* origin. *Molecular and General Genetics* **232**, 460–469.
- Van de Rhee, M.D., Lemmers, R. and Bol, J.F.** (1993) Analysis of regulatory elements involved in stress-induced and organ-specific expression of tobacco acidic and basic β -1,3-glucanase genes. *Plant Molecular Biology* **21**, 451–461.
- Van Kan, J.A.L., Joosten, M.H.A.J., Wagemakers, C.A.M., van den Berg-Velthuis, G.C.M. and de Wit, P.J.G.M.** (1992) Differential accumulation of mRNAs encoding extracellular and intracellular PR proteins in tomato induced by virulent and avirulent races of *Cladosporium fulvum*. *Plant Molecular Biology* **20**, 513–527.
- Vögeli-Lange, R., Fründt, C., Hart, C.M., Nagy, F. and Meins, F.** (1994) Developmental, hormonal, and pathogenesis-related regulation of the tobacco class I β -1,3-glucanase B promoter. *Plant Molecular Biology* **25**, 299–311.
- Watkins, J.T. and Cantliffe, D.J.** (1983) Mechanical resistance of the seed coat and endosperm during germination of *Capsicum annuum* at low temperature. *Plant Physiology* **72**, 146–150.

Watkins, J.T., Cantliffe, D.J., Huber, D.J. and Nell, T.A. (1985) Gibberellic acid stimulated degradation of endosperm in pepper. *Journal of the American Society for Horticultural Science* **110**, 61–65.

Wu, C.-T. and Bradford, K.J. (2002) Class I chitinase is expressed specifically in the micropylar region of *gib-1* tomato seeds in response to wounding or methyl jasmonate. *Abstract of the Seventh International Workshop on Seeds, Salamanca, Spain*.

Wu, C.-T., Leubner-Metzger, G., Meins, F. and Bradford, K.J. (2000) Class I β -1,3-glucanase and chitinase are expressed in the micropylar endosperm of tomato seeds prior to radicle emergence. *Plant Physiology* **126**, 1299–1313.

Received 10 July 2002
accepted after revision 8 January 2003
© CAB International 2003